

# Neutron halo, tritons and alpha clusters in the nucleus ${}^8\text{He}$

**Afsar Abbas**

Institute of Physics  
Bhubaneswar-751005, India  
email: afsar@iopb.res.in

## Abstract

Recent experiments clearly demonstrate the importance of the excited state  $2^+$  at 1.8 MeV in the substructure  ${}^6\text{He}$  for the ground state of the nucleus  ${}^8\text{He}$ . It is shown here that this means that the ground state structure of  ${}^8\text{He}$  rather than being  $\alpha + 4n$  is actually  ${}^3\text{H} + {}^3\text{H} + 2n$ . As such the pair of tritons would form a compact core around which the two valence neutrons provide a halo structure in  ${}^8\text{He}$ . Hence both  ${}^6\text{He}$  and  ${}^8\text{He}$  are two neutron halo nuclei but with cores of quite different nature. All this demonstrates significance of triton clustering in neutron rich nuclei.

The nuclei  ${}^6\text{He}$  and  ${}^8\text{He}$  are uniquely basic for the study of light neutron rich nuclei. While the nucleus  ${}^6\text{He}$  seems to have a structure  $\alpha + 2n$ , with the two neutrons making up a clear cut halo structure around an inert  $\alpha$  [1], the situation for  ${}^8\text{He}$  is known to be more complicated [1]. Though there does seem to be a halo structure in  ${}^8\text{He}$ , however one is not clear as to the nature of the same. Hence it is important to know as to what exactly is going on in  ${}^8\text{He}$ .

In focus here would be two very recent experiments which give interesting insights about the structure of  ${}^8\text{He}$ . These two very different and independent experiments [2,3] reveal identity of structure for  ${}^8\text{He}$ . These seem to be ruling out the standard and canonical ( but somewhat simplistic )  $\alpha + 4n$  structure of  ${}^8\text{He}$ . Hence as such these two experiments cannot be understood in terms of the conventional model of this nucleus. Here we shall discuss how a recent model proposed by the author [4] is able to provide a consistent and revealing understanding of the puzzles provided by these two experiments [2,3].

The first experiment [2] is that of the study of scattering of  ${}^8\text{He}$  on  ${}^4\text{He}$  with the aim of detecting the  $4n$  structure of  ${}^8\text{He}$ . There have been some recent claims of observation of tetra-neutrons in light nuclei. This particular study was done with the aim of understanding this putative tetra-neutronic structure in  ${}^8\text{He}$ . They start with the standard assumption that  $\alpha$  -particles form a well defined and inert object in  ${}^8\text{He}$ . They considered a one step direct  $4n$  transfer process and a two step sequential transfer process of  $2n$  clusters proceeding through the intermediate ground state and the first excited  $2^+$  state ( at 1.8 MeV ) in  ${}^6\text{He}$ . Their DWBA calculations indicated that the two step  $2n$  transfer is clearly more important than the one step  $4n$  transfer. This experiment therefore clearly disfavors the  $\alpha + 4n$  cluster structure of  ${}^8\text{He}$  [2].

In the second experiment [3] they studied the two neutron transfer reactions  $p({}^8\text{He}, t){}^6\text{He}_{gs}$  and  $p({}^8\text{He}, t){}^6\text{He}^*(2^+)$  for the ground state and the first excited states of  ${}^6\text{He}$  respectively. One expects that if the ground state of  ${}^8\text{He}$  contains the subsystem  ${}^6\text{He}$  mainly in the  $0^+$  state, then this would be preferentially populated in  ${}^8\text{He}(p, t)$  reaction. Then it being a second order process, the cross section for the population of  ${}^6\text{He}^*(2^+)$  would be considerably lower. Contrary to these expectations they found the cross section of the  $p({}^8\text{He}, d){}^6\text{He}^*(2^+)$  reaction to be much higher than that of  $p({}^8\text{He}, d){}^6\text{He}_{gs}$ . One therefore concludes [3] that the ground state of  ${}^8\text{He}$

contains subsystem  ${}^6\text{He}$  in the excited state  $2^+$  with a large weight.

Hence what both these two experiments are showing is that  ${}^8\text{He}$  has a two-tier structure with its ground state having  ${}^6\text{He}$  in an the excited state ( $2^+$  at 1.8 MeV ). Therefore the structure of  ${}^8\text{He}$  is very different from that of  ${}^6\text{He}$  which is two loosely bound neutrons outside an inert  $\alpha$ . But here too though  ${}^6\text{He}$  is in an excited state still it is commonly believed that the  $\alpha$  inside it remains inert [1,3].

However this view ignores the fact that in the excited state  $2^+$  the structure of  ${}^6\text{He}$  may not remain pure  $\alpha + 2n$  but may go over to another configuration - that of  ${}^3\text{H} + {}^3\text{H}$  ( ie t+t ). In the past there have been several attempts to understand the structure of  ${}^6\text{He}$  as consisting both of  $\alpha + 2n$  and of t+t cluster structures ( see [7] for references about this ). So one may justifiably ask if it could be that it is the t+t structure of  ${}^6\text{He}$  that is manifesting itself in this excited state relevant to  ${}^8\text{He}$ . And this is exactly what my model predicts [4]. To understand as to how this is possible, first a few words about the model.

Recently using QCD and quark model based ideas, the author has pointed out common threads in such diverse nuclear phenomenon as [4]: 1. The halo phenomena is neutron rich nuclei, 2. The formation and persistence of clusters in nuclei and 3. The nuclear molecule effects. 4. The hole at the centre of the density distribution in  ${}^3\text{H}$ ,  ${}^3\text{He}$  and  ${}^4\text{He}$  [5,6].

It was pointed out by the author that all these require an understanding of two or more nucleons strongly overlapping over a small local region of size  $\leq 1\text{fm}$  in some specific nuclei. This necessarily requires considerations of multi-quark configurations like 6-, 9- and 12-quarks. It was shown by the author [4,5,6] that the nucleons do not like to go into these configurations and thereby the above mentioned nuclear effects are given a consistent explanation. The reader may refer to [4,5,6] for further details.

Let us here concentrate upon a very significant empirical prediction of this new model. That is the effect of triton clustering for neutron rich nuclei. It was shown that as more and more neutrons are added to light nuclei there is a marked effect that an  $\alpha$  plus two neutrons prefer to go to a configuration of two tritons. There is a continuous competition between an  $\alpha$  cluster plus two neutrons remaining as such versus these changing into two clusters of tritons. For a single such configuration in  ${}^6\text{He}$  this breakup does not happen and to a very good approximation  ${}^6\text{He}$  remains an  $\alpha$  plus two neutrons for the ground state [1]. Also consequences like formation of nuclear molecular

structures [4] and also providing us with a better insight into the structure of other  $A=6$  nuclei like  ${}^6\text{Li}$  and  ${}^6\text{Be}$  [7]. The effect of triton ( as well as  $\alpha$  ) clustering also helps us to explain the phenomena of neutron halos [4] and provides a complete and consistent description of all nuclei known to have neutron halo structure [4,8]. It also predicts as to where to look for further new neutron halo nuclei,

People have always thought that for stable  $N=Z$  nuclei with  $4n$  the alphas give a good cluster structure because of the high value of binding energy of this particular nucleus. But the author pointed out [4] that other properties which are significant in this respect are the extremely high surface density of  $\alpha$  and also that it has a hole at the centre. What the other  $A=3$  nuclei,  ${}^3\text{He}$  and  ${}^3\text{H}$  have in common with  $\alpha$  are having holes in the centre and having very high densities on the surface - much higher than any other nuclei [4,5,6]. It is these two properties which all these three light nuclei (  $A=4$  and  $A=3$  ) have in common which makes them ideal candidates around which cluster structures may develop [4]. This is the reason why the tritons are showing up so prominently in forming clusters in neutron rich nuclei. Also for the same reason  ${}^3\text{He}$  would play a similar role in proton rich nuclei [4].

This tendency for neutron rich nuclei to form tritons was used by the author [4] to predict that  ${}^9\text{Li}$  can be treated as made up of  $3t$  and thereby forming a more compact nucleus. Thus when one goes to  ${}^{11}\text{Li}$  one has two loosely bound neutrons forming a loose halo around a compact core of  ${}^9\text{Li}$ . However for  ${}^6\text{He}$  the core  $\alpha$  remains inert and it is the two valence neutrons which form the halo. But when one adds two more neutrons around  ${}^6\text{He}$  to form  ${}^8\text{He}$  then for the ground state of this nucleus there is a distinct change in the substructure of  ${}^6\text{He}$  as discussed above. As per our model discussed here, The excited  $2^+$  state of it can be imagined to have gone into the predominant  $t+t$  configuration. This forms a very compact structure as its two constituent, the tritons themselves are so compact. Therefore it would have high surface densities and thereby preventing the two valence neutrons around it ( for  ${}^8\text{He}$  ) from going through and thus forming a halo around it. Hence as per our model  ${}^8\text{He}$  is also a two neutron halo nucleus with a compact core of two tritons.

In trying to understand the surprisingly large branching ratio for beta-delayed triton emission in  ${}^8\text{He}$ , the authors of ref. [9] suggested the existence of a new  $1^+$  state at 9.3 MeV in  ${}^8\text{Li}$ . They did not know the origin of this state though. Quite clearly in our model here this is the state built up on

the  $1^+$  state at 5.65 MeV in  ${}^6\text{Li}$  ( see [7] for details ). For our purpose here just note that this excited  $1^+$  state in  ${}^8\text{Li}$  has the cluster structure  $(\alpha\text{-t-n})$  [7]. Therefore tritons would be easily emitted in this branch of beta decay in  ${}^8\text{He}$  whose structure as shown here is  $\text{t+t+2n}$ . The beta transition would connect these two structures in a naturally simple manner.

In summary, two recent experimental results are used here to give a new understanding of the structure of the nucleus  ${}^8\text{He}$ . Author's recent model, which has been successfully applied to provide a comprehensive perspective of several diverse nuclear phenomena, again with the insights obtained from these new experiments provides a new description of  ${}^8\text{He}$ . It is predicted to have a structure of  $\text{t+t+2n}$ . Here the two tritons form a compact core ( as an excited state of  ${}^6\text{He}$  ) around which the two valence neutrons form a loose halo.

## References

1. I.Tanihata, D. Hirata, T. Kobayashi, S. Shimoura, K. Sugimoto and H. Toki, Phys. Lett. **B289** (1992) 261
2. R. Wolski, S. I. Sidorchuk, G. M. Ter-Akopian, A. S. Fomichev, A. M. Rodin, S. V. Stepantsov, W. Mittig, P. Roussel-Chomoz, H. Savajols, N. Alamanos, F. Auger, V. Lapoux, R. Raabe, Yu. M. Tchuvil'sky and K. Rusek, Nucl. Phys. **A 722** (2003) 55c
3. A. A. Korshennikov, Nucl. Phys. **A 722** (2003) 157c
4. A. Abbas, Mod. Phys. Lett. **A16** (2001) 755.
5. A. Abbas, Phys. Lett. **B167** (1986) 150.
6. A. Abbas, Prog. Part. Nucl. Phys. **20** (1988) 181.
7. A. Abbas, "Structure of A=6 nuclei:  ${}^6He$ ,  ${}^6Li$  and  ${}^6Be$ ", IOP Preprint:IP/BBSR/2003-18(June'03): physics/0306186
8. A. Abbas, "Triton clustering in neutron rich nuclei and Ikeda-like diagrams", IOP Preprint:IP/BBSR/2003-25(July'03):physics/0307066
9. M.J.G. Borge, L. Johannsen, B. Jonson, T. Nilsson, G. Nyman, K. Riisager, O. Tengblad and K. Wilhelmsen Rolander, Nucl. Phys. **A560** (1993) 664.